

Thermodynamic approach to dense granular matter: a numerical realization of a decisive experiment.

Hernán A. Makse¹, and Jorge Kurchan²

¹ *Levich Institute and Physics Department, City College of New York, New York, NY 10031, US*

² *P.M.M.H., Ecole Supérieure de Physique et Chimie Industrielles, 10 rue Vauquelin 75231 Paris, France.*

(February 1, 2008)

Years ago Edwards proposed a thermodynamic description of dense granular matter, in which the grains (the ‘atoms’ of the system) interact with inelastic forces. The approach is intriguing but is not justified from first principles, and hence, in the absence of conclusive tests of its validity, it has not been widely accepted. We perform a numerical experiment with a realistic granular matter model specially conceived to be reproducible in the laboratory. The results strongly support the thermodynamic picture.

The similarities between a driven granular system and a system of molecules in a fluid have prompted the use of statistical ideas to describe granular matter [1], such as for example kinetic theories of gases to study rapid flows of low-density systems of inelastic particles [2]. In this context, it is customary to associate the mean kinetic energy of the particles with a ‘granular temperature’, having no real thermodynamic meaning.

For the opposite limit of dense, slowly flowing granular matter, a more radical train of ideas was initiated more than a decade ago by Edwards and coworkers: the proposal of a statistical ensemble [3–6], and through it, thermodynamic notions such as entropy and temperature (the latter unrelated to the ‘granular temperature’ above). Although the idea of a thermodynamic description of granular matter was recognized as attractive, it was not universally accepted because there is no known first principle justification of Edwards’ statistical ensemble, such as there is for ordinary statistical mechanics of liquids or gases (Liouville’s theorem).

Recently, analytical developments and numerical work on schematic models originally devised for glasses have given a new perspective for the dynamics of dense granular matter [7–12]. The theoretical support provided by this framework for a statistical approach has spurred a renewed interest in Edwards’ thermodynamics.

On the experimental side, years ago Nowak *et al.* [13] studied in detail the density fluctuations in a vibrated granular material, and proposed that these fluctuations should reflect an underlying thermodynamics, much as they do in an ordinary thermal system. However, as far as the actual verification of Edwards’ approach, the evidence they found was if anything rather negative: we shall discuss below some possible reasons for this. On the other hand, Edwards’ approach has never been tested with simulations of realistic models of granular materials, characterized by the fact that energy is supplied by external driving (via shear or tapping) and dissipated by inelastic collisions and slippage between the grains.

In this work we describe a numerical experiment specially conceived to be reproducible in the laboratory, using a realistic model of sheared granular matter. Con-

sidering particles of different sizes inside the medium, we compute a dynamical temperature from an Einstein relation relating random diffusion and mobility (transverse to the shear direction) at long time scales. If there is an underlying thermodynamics, it will impose that the dynamical temperature be independent of the tracer’s shape: a very strong condition which we show to hold. We then perform an explicit computation to show that the temperature arising from Edwards’ thermodynamic description and the dynamic one measured from Einstein’s relation coincide. This last step cannot be performed in the laboratory, so the numerical simulation provides the missing link between thermodynamic ideas and diffusion-mobility checks.

Consider a ‘tracer’ body of arbitrary shape immersed in a liquid in equilibrium at temperature T . As a consequence of the irregular bombardment by the particles of the surrounding liquid, the tracer performs a diffusive, fluctuating ‘Brownian’ motion. The motion is unbiased, and for large times the average square of the displacement goes as $\langle |x(t) - x(0)|^2 \rangle = 2Dt$, where D is the diffusivity. On the other hand, if we pull gently from the tracer with a constant force f , the liquid responds with a viscous, dissipative force. The averaged displacement after a large time is $\langle [x(t) - x(0)] \rangle = f\chi t$, where χ is the mobility. Clearly, the same liquid molecules are responsible both for the fluctuations and for the dissipation. Although both D and χ strongly depend on the shape and size of the tracer, they turn out to be always related by the Einstein relation $D/\chi = T$, where T is the temperature of the liquid.

Conversely, if in a fluid of unknown properties we find that several tracers having different diffusivities and mobilities yield the same ratio $D/\chi = T$, we may take this as a strong evidence for thermalisation at temperature T . Indeed, recent analytic schemes for out of equilibrium glassy dynamics have suggested the existence of such a temperature T governing the slow components of fluctuations and responses of all observables [7,10].

In order to test the existence of this temperature for dense granular matter, we perform, with a realistic numerical model, a diffusion-mobility experiment in con-

ditions that can be reproduced in the laboratory. In the model, deformable spherical grains interact with one another via non-linear elastic Hertz normal forces, and non-linear elastic and path-dependent Mindlin transverse forces [14]. The normal force, F_n , has the typical $3/2$ power law dependence on the overlap between two spheres in contact, while the transverse force, F_t , depends on both the shear and normal displacements between the spheres. For two spherical grains with radii R_1 and R_2 : $F_n = \frac{2}{3} k_n R^{1/2} w^{3/2}$, $\Delta F_t = k_t (Rw)^{1/2} \Delta s$. Here $R = 2R_1 R_2 / (R_1 + R_2)$, the normal overlap is $w = (1/2)[(R_1 + R_2) - |\vec{x}_1 - \vec{x}_2|] > 0$, and $\vec{x}_1, \vec{x}_2$ are the positions of the grain centers. The normal force acts only in compression, $F_n = 0$ when $w < 0$. The variable s is defined such that the relative shear displacement between the two grain centers is $2s$. The prefactors $k_n = 4G/(1 - \nu)$ and $k_t = 8G/(2 - \nu)$ are defined in terms of the shear modulus G and the Poisson's ratio ν of the material from which the grains are made (typically $G = 29$ GPa and $\nu = 0.2$, for spherical glass beads). We assume a distribution of grain radii in which $R_1 = 0.105$ mm for half the grains and $R_2 = 0.075$ mm for the other half. The observables are measured in reduced units: length in units of R , force in units of GR^2 , time in units of $\sqrt{\rho R^2/G}$, where ρ is the density of the particles. Internal dissipation is included in two ways: (1) via a viscous damping term proportional to the relative normal and tangential velocities, and (2) via sliding friction: when F_t exceeds the Coulomb threshold, μF_n , the grains slide and $F_t = \mu F_n$, where μ is the friction coefficient between the spheres (typically $\mu = 0.3$).

We perform molecular dynamics (MD) simulations for a binary system of 200 large and small spheres in a periodic 3D cell. Our calculation begins with a numerical protocol designed to mimic the experimental procedure used to prepare dense packed granular materials. The simulations start with a gas of spherical particles located at random positions in a periodically repeated cubic cell of side L . The system is then compressed and extended slowly until a stationary situation with a specified value of the pressure and volume fraction—above the random close packing fraction—is achieved. We then apply a gentle shear in the y -direction at constant volume by moving the periodic images at the top and bottom of the cell with velocities $\dot{\gamma}L/2$, where $\dot{\gamma}$ is the shear rate (Lees-Edwards boundary conditions). In this simple shear flow the gradient of the velocity along z is uniform (no shear bands). We apply a shear rate of $\dot{\gamma} = 10^{-4}$, which allows the system to be in the quasi-static regime. We focus our study on the slow shear rate, quasi-static limit where the system is always close to jamming, but moves just barely enough to avoid stick-slip motion [15]. Here the external pressure is large enough (~ 10 MPa in a typical simulation) and the average coordination number is high enough (typically ~ 7) that deformation and elasticity of the particles play the dominant role, as opposed to

the collision dominated rapid flow regime described by kinetic theories. We checked that shear induced segregation is absent at the times scales of our simulations. The contacts between particles are enduring and the internal stresses in the system are transmitted via a network of ‘force chains’ [16,1], independent of the shear rate.

After a transient of the order of the inverse shear rate, we start measuring the spontaneous $\langle |x(t) - x(0)|^2 \rangle$ and force-induced displacements $\langle [x(t) - x(0)]/f \rangle$ along the x -direction for the two types of particles with different sizes. The results are shown in Fig. 1. We notice that the diffusivities and the mobilities are different for the two type of particles, since they have different sizes. However, when we draw the parametric plot of $\langle |x(t) - x(0)|^2 \rangle$ versus $\langle [x(t) - x(0)]/f \rangle$ (Fig. 2) we find parallel straight lines for large time scales, implying an extended Einstein relation:

$$\langle |x(t) - x(t_w)|^2 \rangle = 2 T_{dyn} \frac{\langle [x(t) - x(t_w)] \rangle}{f}, \quad (1)$$

valid for both particles *with the same* T_{dyn} for widely separated time scales $t \gg t_w$. This suggests that T_{dyn} can be considered to be the temperature of the slow modes.

For small time scales (fast rearrangements, near the origin in Fig. 1) the fluctuation-dissipation plot shows that there is not a well defined Einstein-relation temperature. We expect this result since the fast motion of the grains depends on the microscopic interactions dominated by inelastic collisions [17]. We also calculate the ‘granular kinetic temperature’ defined in terms of the velocity fluctuations in the x -direction. Unlike T_{dyn} , we find that this temperature is different for the two types of particles, and their values are two orders of magnitude smaller than T_{dyn} . We expect this result since the kinetic granular temperature is dominated by the fast (high frequency) modes, which are not thermalised to T_{dyn} .

We also repeat the numerical experiment for a system of Hertz spheres without transverse forces ($\mu = 0$: experimental realizations of elastic spheres with viscous forces but without sliding friction are foams and compressed emulsions [18,19]) and find that T_{dyn} is well defined at long time scales for this case as well (see Fig. 1). Thus, our results suggest that the validity of a structural temperature for long-scale displacements (larger than a fraction of the particle size) holds in the presence of viscous forces between grains or even of a sliding threshold (Coulomb's law).

The existence of a single temperature is an instance of the zero-th law of thermodynamics, for which we find positive evidence here, at variance with the experimental result in Nowak *et al* [13]. Three possible reasons for the apparent violation of the ‘zero-th law’ in their experiments are *i*) the effect of an unknown height-dependent pressure (as pointed out by authors), *ii*) a rather high tapping amplitude (Edwards ensemble is in principle only valid in the high compaction limit [10]) and *iii*) the fact

that the density fluctuations considered are integrated over all frequencies, thus including also ‘fast’ relaxations.

Next, we treat the question whether it is possible to relate the dynamical temperature obtained above to the thermodynamic construction of slowly driven out of equilibrium systems proposed by Edwards. Whereas in the Gibbs construction for equilibrium statistical mechanics one assumes that the physical quantities are obtained as average over all possible configurations, Edwards ensemble consists of only the blocked configurations (static or jammed) at the appropriate energy and volume. The strong ‘ergodic’ hypothesis is that all blocked configurations of given volume and energy can be taken to have equal statistical weights. This formulation leads to an entropy $S_{Edw}(E, V)$, and the corresponding temperature $T_{Edw}^{-1} = \partial S_{Edw} / \partial E$ and compactivity $X_{Edw}^{-1} = \partial S_{Edw} / \partial V$.

In order to calculate T_{Edw} and compare with the obtained T_{dyn} we need to count the number of blocked configurations at a given energy and volume. (For this calculation we concentrate in the case without tangential forces and sliding friction, in order to avoid path dependency which would lead to an ambiguity in the definition of blocked configurations—see below). To do this in practice, we extend the ‘auxiliary model’ method [10] to the case of deformable grains. We define an auxiliary model composed of the true deformation energy (E , the Hertzian energy of deformation of the particles) and of a term that vanishes in (and only in) the blocked configurations: $E_{block} \propto \sum_a |\vec{F}_a|^2$. (Here $\vec{F}_a$ is the total contact force exerted on particle a by its neighbours). We perform equilibrium MD with two auxiliary temperatures (T^*, T_{aux}), corresponding to the partition function $\sum \exp[-E/T^* - E_{block}/T_{aux}]$. Annealing T_{aux} to zero selects the blocked configurations ($E_{block} = 0$), while T^* fixes the energy E . We start by equilibrating the system at high temperatures (T_{aux} and $T^* \sim \infty$) and anneal slowly the value T_{aux} to zero and tune T^* so as to reach the observed value of E during shear. At the end of the process ($T_{aux} \rightarrow 0$), $T^*(E) = T_{Edw}(E)$, since in this limit we are sampling the configurations with vanishing fraction of moving particles at a given E .

Figure 3 shows various annealing protocols. We plot the value of the true deformation energy as a function of T_{aux} during annealing using different values of T^* as indicated in the figure. At the end of the annealing process the system stabilizes at a given deformation energy. Only when T^* is equal to T_{dyn} obtained through the Einstein relation during shear, we find that the final E coincides, within the accuracy of the simulations, with the mean energy of the system under shear. For other values of T^* the annealed system equilibrates at energies which are above of below the distribution of deformation energies during shear as seen in Fig. 3. *This proves the equality between Edwards’ and the dynamic temperature.*

We conclude with some remarks: *i)* Since the blocked configurations are the same whatever the inter-grain dissipation coefficient, Edwards’ ensemble (and hence its temperature) are *insensitive to viscous dissipation*.

ii) On the contrary, tangential forces and sliding friction block certain configurations, depending on how they are accessed. The ensemble of blocked configurations is then ill-defined. We have not tried to construct any ensemble, but content ourselves with the observation that T_{dyn} is also in this case independent of the particle size (Fig. 2)—we suspect that thermodynamic concepts apply, but that the relevant ensemble goes beyond Edwards’ construction as it stands.

iii) We have tested the validity of the thermodynamics in an ideal homogeneous system with periodic boundary conditions by explicitly avoiding structural features of dense granular flows such as shear bands and segregation of the species. Even though it remains to be seen whether the thermodynamic picture can account for these inhomogeneous effects, our ideal system may prove to be useful in deriving constitutive relations to be used in macroscopic theories of slow granular flows.

To summarize: we have performed numerically an experiment with dense granular systems specially conceived to be a ‘dress rehearsal’ for the real laboratory one. The independence of the Einstein-relation temperature on the tracer provides a strong test for the thermodynamic ideas. We have also showed that this temperature, obtained from measurable quantities, indeed matches Edwards’ one. If, as now seems likely, this result is confirmed in the laboratory, this will give an experimental foundation for the use of the powerful tools of statistical mechanics as the framework to study this kind of far-from-equilibrium dissipative dynamical system.

-
- [1] H. M. Jaeger, S. R. Nagel, R. P. Behringer, *Rev. Mod. Phys.* **68**, 1259 (1996).
 - [2] J. T. Jenkins, S. B. Savage, *J. Fluid Mech.* **130**, 187 (1983).
 - [3] S. F. Edwards, in *Granular Matter: An Interdisciplinary Approach*, A. Mehta, Ed., (Springer-Verlag, New York, 1994), pp. 121-140, and references therein.
 - [4] S. F. Edwards, in *Disorder in Condensed Matter Physics*, Blackman and Taguena, Eds., (Oxford University Press, Oxford, 1991), pp. 147-154.
 - [5] A. Mehta, R. J. Needs, S. Dattagupta, *J. Stat. Phys.* **68**, 1131 (1992).
 - [6] S. F. Edwards and D. V. Grinev, *Phys. Rev. E* **58** (1998) 4758.
 - [7] L. F. Cugliandolo, J. Kurchan, L. Peliti, *Phys. Rev. E* **55**, 3898 (1997).
 - [8] M. Nicodemi, *Phys. Rev. Lett.* **82**, 3734-3737 (1999).

- [9] Kurchan J., in ‘Jamming and Rheology: Constrained Dynamics on Microscopic and Macroscopic Scales’ (1997), <http://www.itp.ucsb.edu/online/jamming2/>; cond-mat 9812347, S.F., Liu, A. and Nagel, S.R. Eds., Taylor and Francis, New York (2001).
- [10] For a background on the connections with dynamical glass theory, and an implementation on schematic finite dimensional models, see: A. Barrat, J. Kurchan, V. Loreto, M. Sellitto, *Phys. Rev. Lett.* **85**, 5034 (2000); *Phys. Rev. E* **63** (2001) 51301, and references therein.
- [11] J. Javier Brey, A. Prados, B. Sanchez-Rey, *Physica A* **275**, 310 (2000).
- [12] A. Lefèvre, D. S. Dean, *Eur. Phys. Journal* **B21** (2001) 121, and cond-mat/0106220.
- [13] E. R. Nowak, J. B. Knight, E. Ben-Naim, H. M. Jaeger and S.R. Nagel, *Phys. Rev. E* **57** (1998) 1971.
- [14] K. L. Johnson, *Contact Mechanics* (Cambridge University Press, Cambridge, 1985).
- [15] S. B. Savage, *Adv. Appl. Mech.* **24**, 289 (1994).
- [16] Dantu, P “A Contribution to the Mechanical and Geometrical Study of Non-cohesive Masses”, in *Proc. 4th Int. Conf. on Soil Mechanics Foundation Engineering*, pp. 144 (Butterworths, London, 1957).
- [17] This difference between fast and slow relaxations is similar to the one found for glasses, although in the case of glasses the fast modes are equilibrated at the bath temperature [7].
- [18] D. J. Durian, *Phys. Rev. Lett.* **75**, 4780 (1995).
- [19] M.-D. Lacasse, G. S. Grest, D. Levine, T. G. Mason, D. A. Weitz, *Phys. Rev. Lett.* **76**, 3448 (1996).




FIG. 2. Parametric plot of diffusion vs response function for small and large grains and for spheres interacting with tangential forces and without tangential forces (Coulomb frictionless). The fitting at long time scales shows the existence of a well defined temperature which is the same for small and large grains: $T_{dyn} = 2.8 \times 10^{-5}$ for grains without transverse forces and $T_{dyn} = 1.2 \times 10^{-4}$ for grains with Mindlin transverse forces and Coulomb friction. We calculate the response function for several small external fields and find the same temperature indicating that we are in the linear response regime. Plotted are results for a system without transverse forces using: $f = 1.7 \times 10^{-5}$ (small grains $\square$, large grains $\circ$) and $f = 2.6 \times 10^{-5}$ (small grains $\blacksquare$, large grains $\bullet$). For a system with tangential forces and Coulomb friction we show the case $f = 6 \times 10^{-5}$ (small grains $\triangle$, large grains $\times$).




FIG. 1. (A) Diffusion and (B) response function for the large and small particles in a sheared granular material measured perpendicular to the shear plane as a function of time in MD steps. Both quantities depend linearly on time at the late stages of the evolution. The response function is measured by applying a constant force f in the x -direction to each type of particle. Averages are taken over 30 temporal realizations. The obtained diffusivities and mobilities depend on the particles size as expected.




FIG. 3. Annealing procedure to calculate T_{Edw} at different true energies. We plot the true deformation energy versus T_{aux} together with the distribution of deformation energies obtained during shear (dashed curve, mean value $\langle E \rangle = 8.4 \times 10^{-4}$). We equilibrate the system for 40 million iterations at A: ($T^* = 3.4 \times 10^{-2}$, $T_{aux} = 3 \times 10^{-8}$). We then anneal slowly both temperatures until B: ($T^* = 3.4 \times 10^{-4}$, $T_{aux} = 3 \times 10^{-10}$), where we split the trajectory in three paths in the (T^*, T_{aux}) plane. Path 1: we anneal $T_{aux} \rightarrow 0$ and $T^* \rightarrow 2.8 \times 10^{-5}$ which corresponds to T_{dyn} obtained during shear (Fig. 2). Path 2: we anneal $T_{aux} \rightarrow 0$ and $T^* \rightarrow 3.4 \times 10^{-6}$. Path 3: we anneal $T_{aux} \rightarrow 0$ but keep $T^* = 3.4 \times 10^{-4}$ constant. When we set $T^* = T_{dyn}$ (Path 1), the final true energy value when $T_{aux} \rightarrow 0$ falls inside the distribution of energies obtained during shear and it is very close to $\langle E \rangle$. This proves that $T_{dyn} = T_{Edw}$ under the numerical accuracy of the simulations. For other values of $T^* \neq T_{dyn}$ the final E falls out of the distribution obtained during shear (Paths 2 and 3). We also follow different trajectories (not shown in the figure) to $T^* \rightarrow 2.8 \times 10^{-5}$, $T_{aux} \rightarrow 0$ and find the same results indicating that our procedure is independent of the annealing path.